

First record of *Lernaea cyprinacea* (Linnaeus, 1758) (Copepoda: Cyclopoida) on *Betta splendens* in Brazil

Elton Lima Santo^[1], Themis de Jesus Silva^[2], Misleni Ricarte de Lima^[3],
Ricardo Fábio Teodósio Cavalcante Junior^[4], Sarah Jacqueline Cavalcanti da Silva^[5],
Emerson Carlos Soares^[6]

[1]elton.santos@ceca.ufal.br. [2]themisjdasilva@gmail.com. [3]misleniricarte@hotmail.com. [4]ricardo.ftjr@gmail.com.
[5]sarah.silva@ceca.ufal.br. [6]soaemerson@gmail.com. Campus de Engenharia e Ciências Agrárias, Universidade Federal de Alagoas.

ABSTRACT

The aim of this work is to bring this first record of *L. cyprinacea* in *B. splendens*. The anchor worm, also known as *L. cyprinacea* is an ectoparasitic copepod that can result in the mortality of the host by causing hemorrhages, ulcerations, and secondary infections. *L. cyprinacea* is widely distributed in world. Previous reports are restricted cyprinaceae on other fishes. The study detected, for the first time, the occurrence of lernaecosis in *Betta splendens* of the Alagoas state, Brazil. The occurrence was *L. cyprinacea* was detected on the narines, gills, soft parts the dorsal and anal fin of *Betta splendens*. It brings concern the occurrence of the anchor worms *Lernaea cyprinacea* in a cultivate because this exotic species is able to spread and to parasitize other species of fishes.

Keywords: Anchor worms. Copepod. ornamental fish. ectoparasite.

RESUMO

O objetivo deste trabalho é trazer este primeiro registro de *L. cyprinacea* em *B. splendens*. O *L. cyprinacea*, também conhecido como: verme âncora, é um copépode ectoparasitário que pode causar mortalidade do hospedeiro, causando hemorragias, ulcerações e infecções secundárias. O *L. cyprinacea* é amplamente distribuído no mundo. Relatos anteriores são restritos a cyprinaceae em outros peixes. O presente estudo encontrou pela primeira vez, a ocorrência de lernaecose em *Betta splendens*, provinda do Estado de Alagoas, Brasil. A ocorrência de *L. cyprinacea* foi detectada nas narinas, brânquias, partes moles da barbatana dorsal e anal de *Betta splendens*. Preocupa a ocorrência do verme âncora, *Lernaea cyprinacea*, em um cultivo, porque esta espécie exótica é capaz de se espalhar e parasitar outras espécies de peixes.

Palavras-chave: vermes âncora. copepodo. peixe ornamental. ectoparasita.

1 Introduction

With the rapid developments of commercial aquaculture farming, diseases to become the major constraint of the development of aquaculture globally. According Santos *et al.* (2013) most of the pathogens are opportunist to infect cultured fish once the environment is deteriorated. Parasitic diseases constitute impact significant of diseases affecting warm water fish in world, that mainly affect the fish health, growth and survivability (SALINAS *et al.*, 2019).

In recent decades, the relevance of studies related to parasites and other pathogens of aquatic organisms, especially those hosts with potential for cultivation and marketing, given the significant increase in these activities in Brazil and worldwide (LUQUE, 2004; SANTOS *et al.*, 2017).

Lernaea cyprinacea (Linnaeus, 1758) is among most known species of the copepods exotic that parasite fish in Brazil (TAVARES-DIAS *et al.*, 2015). The parasite *L. cyprinacea* and most exotic species have been a worldwide environmental concern.

Introduced during the 80's via the common carp *Cyprinus carpio* Linnaeus, 1758, *L. cyprinacea* difusion began at the northeast area, spreading to southeastern, mid-west and reaching the all regions of Brazil (FORTES; HOFFMANN; SCARIOT, 1998). Its economic importance in brazilian fish farm is increasing due to numerous epizootics occurring among the most important farmed fish species. However, there are no official reports of *L. cyprinacea* induced fish mortalities from Brazil.

Lernaeids occur in freshwater fishes both in natural water systems and in aquaculture environments. They are notorious killers specifically of small fishes, and re the caused of great economic loss. They are suspected of transmitting viroses and/or bacteria which result in secondary infection.

Soares *et al.* (2018) reports that *L. cyprinacea* is one of the most common parasites in wild and aquaculture fishes of Brazil South region and produces serious economic losses to the fish industry. Therefore, the extensive geographical distribution can not be directly attributed to dispersion by natural means. The human activity could have an indirect influence, by the scaering of *L. cyprinacea* when using live specimens of fish as bait in other lakes and rivers.

Lernaea cyprinacea is an opportunistic ectoparasite of many freshwater fish species and non-fish hosts like amphibians. Adult females of *L. cyprinacea* attach

to exposed body surfaces of host fish including the head, dorsal, ventral, and caudal regions, but most frequently to the bases of fins. Their aggregation on body sites induces severe local damage with adverse consequences to host function and survival (BARSON; MULONGA; NHIWATIWA, 2008).

Worldwide, the popularity of fish keeping has grown by 14% annually since the 1970s and currently over 1 billion individual fish are traded internationally each year (MACEDA-VEIGA *et al.*, 2016). The global aquarium fish trade (freshwater and marine ornamental fish plus accessory products) is worth c. US \$15–30 billion (PENNING *et al.*, 2009). Around 500 million fishes are traded annually by 145 countries, of which 80-85% is tropical species (KUMAR *et al.*, 2016).

Several studies have been developed to improve the knowledge on the biology and management during the rearing of ornamental species, such as: feeding management (SANTOS *et al.*, 2014; SANTOS *et al.*, 2019), techniques and system of production (SALES *et al.*, 2016), nutrition (THONGPRAJUKAEW *et al.*, 2011; ZUANON; SALARO; FURUYA, 2011; TORRES-VALENCIA; IMUES-FIGUEROA; BUCHELI-FUELANTALA, 2016), reproduction (GIANNECCHINI; MASSAGO; FERNANDES, 2012), behavior (MACLAREN; WISNIEWSKI; MACLAREN, 2018), laviculture management (FABREGAT *et al.*, 2017), among others.

The number of species for ornamental purposes is high, and there is lack of information about diseases and parasites, even for those produced globally on a large scale. However, despite the variety in fresh water fish species, certain species are extremely popular amongst hobbyists, including the tropical fish *Betta splendens* Regan, 1910)

This fish is an extremely popular ornamental fish among hobby aquarists. It has na interesting behavioral repertoire, particularly where male aggression and territoriality are concerned (PLEEGING; MOONS, 2017).

Betta splendens is an important species in the Brazil and international ornamental fish trade due its coloured patterns and singular behaviour. Reproduction and larviculture are a challenge for many producers due to the lack of scientific information, mainly on the diseases and parasites.

The Siamese fighting fish (*Betta splendens*) is a member of the Labyrinth fish family (Belontiidae). *B. splendens* in Latin refers to brilliant warrior, indeed a suitable name, since males of the colourful species

perform an elaborate aggressive display when provoked (MOTLAGH *et al.*, 2012).

The species natural habitats are shallow freshwater ponds with muddy bottoms or flooded rice paddy fields. This vegetation provides cover against fish-eating birds, like egrets, herons, kingfishers and others predators (BERTUCCI; MATOS; DABELSTEEN, 2013). They can breathe oxygen from the surface (due to their labyrinth organ), which enables the species to survive in low oxygen waters

The present study reports the presence of *L. cyprinacea* parasitizing *Betta splendens* under captivity in the State of Alagoas, Brazil. A detailed description and previous records in other fishes in the Brazil, are provided.

2 Materials and methods

Our study area is located in the municipality of Maribondo, Alagoas State, Brazil, at local from which *Betta splendens* has been collected (-9.591395 N, -36.307179 E). The local study is characterized by a humid, tropical climate. The maximum temperature is 33 °C with an average minimum 18 °C in month of the July of 2018.

The Fishes, samples of *Betta splendens* (n = 24), only males, were collected of collective field tank of fishes, in Maribondo, Alagoas, Brazil. Posteriorly, examined for the presence of *L. cyprinacea* after collection. Areas around the fins, nostril, operculum and buccal cavity were examined with a dissection microscope. Fishes were put on ice and immediately transported to the laboratory.

To extract crustaceans, fish were subjected to immersion for 10-30 minutes with 1% sodium chloride. Parasites that did not detach on their own were removed with dissecting forceps. Once removed, the affected area was treated with a mild disinfectant.

The ectoparasites were preserved in ethanol 70% for their subsequent identification in the Laboratory of the Aquaculture, Federal University of Alagoas (UFAL, Alagoas, Brasil). The *L. cyprinacea* was identified based on published references following Thatcher (2006).

The review of the hosts and geographical distribution in range of the anchor worm was made from national and international scientific journals. In the case of Brazil, available technical reports, theses, and congress summaries were also included.

Water quality parameters were measured at the surface at each field tank, in original local of animals management. Temperature (°C), nitrite, pH and


dissolved oxygen concentration were determined by using a YSI water meter (Professional Plus).

All experimental procedure followed the Animal Ethics Committee of the Federal University of Alagoas recommendations (n° 07/2018- CEUA/UFAL).


3 Results and discussion

A detailed description of *L. cyprinacea* is presented, including the corresponding measurements (Figure 1 and 2).

Figure 1 – *Betta splendens* parasitized by anchor worm, *Lernaea cyprinacea* (arrows) in gills (a) and dorsal fin (b).



(a)




(b)

In this study, the parasite considered as the cause of lernaecias in ornamental exotic fish species, *Betta splendens*, was *L. cyprinacea*. The morphological characteristics observed here are typical of *L. cyprinacea* (DEMAREE JR, 1967; THATCHER, 2006; KOYUN; ULUPINAR; MART, 2015; HOSSAIN; FERDOUSHI; RUPOM, 2018). It has an elongate, tubular body, which is differentiated into

a cephalothorax, thoracic region and abdomen. The worm's head is distinctly round with antennae and antennules attach laterally on the anterior portion and a posterior anchor processes. Oral appendages and the first pair of swimming legs are situated in the ventral surface of the cephalothorax.

The second, third and fourth pairs of swimming legs are attached to the thorax, whereas the abdômen bears the fifth and sixth pairs of swimming legs. The abdomen is slightly broadened and curved at the posterior region. In similarity that reports from Magalhães (2006) studying *Astyanax bimaculatus* Linnaeus, 1758, in Minas Gerais, Brazil.

Figure 2 – Parasite anchor worm *Lernaea cyprinacea* (a) with eggs (b).



Gervasoni *et al.* (2018) observed the description general morphology, post metamorphic females like: elongated body segmented into cephalothorax, neck and genito-abdomen. In relation to cephalothorax composed of head, mouth structure, four cylindrical projections, lobes or horns and first pair of legs. Corroborating also the present study.

Already, Waicheim *et al.* (2017) related that the specimens of *L. cyprinacea* can be distinguished from

other species of the genus by the following combination of characters: head with 4 large anchors frequently branched, dorsal pair much larger than ventral, dividing into 2 branches some distance from their bases, ventral pair usually simple, neck consisting of second to fourth leg-bearing segments, more or less circular in cross section, and genital pore near posterior extremity, fourth legs on hind body, abdomen conical usually displaced dorsally, and total length 10–20 mm.

Future changes in the aquatic environment may stress fish population even further which may lead to an increase in parasite load in up seven parasites per host) in this species, as Mancini *et al.* (2008) suggested. The current study supports the notion that *L. cyprinacea* can cause death of the hosts, due to intense inflammatory reaction in the subcutaneous tissue as observed. Mirzaei (2015) related that anchor worms can cause intense inflammation, leading to secondary bacterial and fungal infections. These secondary infections sometimes worsen and kill the fish. Larger numbers of parasites on the gill can interfere with respiration, causing death.

Therefore, future studies on parasitic fauna of in ornamental fishes on Brazil are essential to record host and parasite biology and the spread of this pathogen to natural water sources and to other ornamental fishes.

Probably, the infection caused in the bettas were caused by parasites that were parasitizing other fish and that were in the same environment, as observed in *Carassius auratus* and *Poecilia sphenops*, in the same fish farming environment. The fluctuation of temperature may contribute also to the parasitic outbreak.

The parasites can be removed manually by using tweezers as it can be seen with naked eye. However, this method is not practical and unrealistic if fish are heavily infested with huge amount of parasites.

Corroborating this statement, Vasagan *et al.* (2006) report that, *L. cyprinacea* is a common parasite infecting numerous species of freshwater fish and the common method of control is by using removed manually. However, the adult parasite penetrates the fish host and becomes deeply embedded in the tissues making it extremely difficult to remove them (FURTADO *et al.*, 2019).

In the present study, a large parasite infestation in the gills region was verified. Notably harming the health and development of animals. However, *Lernaea cyprinacea* also was detected on the others parts:

narines, soft parts the dorsal and anal fin of *Betta splendens*.

Carriel (2014) reports that gills are of fundamental importance being involved in health maintenance, oxygen uptake and carbon dioxide release, as well as nitrogen waste and osmoregulation processes. Also, the gills are responsible for the basic acid balance and sensory function in tasting. Thus, they have a delicate structure and exposed to environmental changes. Thus, any change in gill structure will affect fish survival and performance. Gill infections cause some localised hyperplasia of the epithelial tissues and also cause intense epithelial proliferation which may seriously interfere with respiration and also support spread of bacterial infection (Shariff, 1986).

It was also observed that parasite aggregations around the eye cause lens destruction and blindness. Innal *et al.* (2017) report that Lernaeid copepods are among the most harmful parasites of freshwater fish, especially young fish, which may be killed by only a few parasites.

Diseased fish not only disturbs the nutrient balance in fish body but also make it unfits nutritionally and pathologically. In Brazil there are few descriptions of parasitic fauna of farmed ornamental fish, as well as the pathogenesis caused by the parasites in their hosts.

In *Betta splendens*, the presence of *Lernaea cyprinacea* has never been reported in the scientific literature. This research is the first report of this parasite in *Betta splendens* in the world. Information such as this is necessary to ensure the production of healthy animals as a subsidy to prevent losses in the activity of ornamental fish farming.

Worth mentioning that, parasites are normally present in the environment or in the body of the fish,

and when there is an imbalance in the environment-host-parasite relationship, outbreaks of disease occur (MARTINS *et al.*, 2002). However, it is known that the practice of introducing exotic species is frequent in aquariums and may be responsible for the spread of parasites, including anchor worm, endangering the entire population of farmed fish.

The average values for temperature, pH, nitrite, and dissolved oxygen during the collection period were 27°C ± 0.5°C; 6.8 ± 0.3; 0.15 mg.L⁻¹ ± 0.05 and 6.50 mg.L⁻¹ ± 0.25, respectively, at each field tank, in original local of animals management. These values remained stable and within the range recommended for the tropical species.

The monitoring of aquatic environments by measuring the density of parasite vectors in the water is very important. Unsustainable aquaculture, commercial and ornamental, has great potential to damage aquatic biodiversity and ecosystem functions and services.

Agostinho; Gomes; Pelicice (2007) account that exotic species can have a negative impact by transmitting diseases or etiological agentes into native communities. Therefore, colonization of exotic pathogens in aquatic systems constitutes a permanente threat to the integrity of natural environments.

This ectoparasite can colonized native fish species and is dangerously invasive when environmental conditions are favorable, as reported in this work. Further research efforts are needed to prevent the role currently played by *L. cyprinacea* in ornamental fish.

The review of the hosts and geographical distribution in range of the anchor worm are expressed in table 1.

Table 1 – Previous records of hosts for *Lernaea cyprinacea* in Brazil.

Province/ State	Location (Original name in portuguese)	Popular and scientific name Hosts	References
Alagoas	municipality of Porto Real do Colégio and Igreja Nova	Tambaqui - <i>Colossoma macropomum</i> Cuvier, 1818	FUJIMOTO <i>et al.</i> (2019)
Rio Grande do Sul	municipality of Antonio Prado	<i>Lambari do rabo amarelo - Astyanax bimaculatus</i> Linnaeus, 1758	GALLIO <i>et al.</i> (2007)
Rio Grande do Sul	Pardinho River	<i>Lambari do rabo vermelho - Astyanax fasciatus</i> Cuvier, 1819	DÜPONT; LOBO (2011)
Rio Grande do Sul	Arroio Creek	<i>Carpa comum - Cyprinus carpio</i> Linnaeus, 1758 <i>Biru - Cyphocharax spilotos</i> Vari, 1987 <i>Biru - Cyphocharax voga</i> Hensel, 1870 <i>Braga - Steindachnerina biornata</i> Braga & Azpelicueta, 1987	QUEROL <i>et al.</i> (2005)

(Continue)

(Continuation)

Province/ State	Location (Original name in portuguese)	Popular and scientific name Hosts	References
Minas Gerais	Paraíba do Sul river	<i>Lambari do rabo amarelo</i> - <i>Astyanax bimaculatus</i>	MAGALHÃES (2006)
Minas Gerais	San Francisco river (barragem três Marias)	<i>Cumbaca</i> - <i>Franciscodoras marmoratus</i> Reinhardt, 1874	SANTOS; BRASIL- SATO (2006)
Paraíba	municipality of Sumé	<i>Traíra</i> - <i>Hoplias malabaricus</i> Bloch, 1794	BERNARDINO et al. (2016)
Paraná	Tibagi river	<i>Pacu</i> - <i>Piaractus mesopotamicus</i> Holmberg, 1886 <i>Carpa comum</i> - <i>Cyprinus carpio</i> Linnaeus, 1758 <i>Piauçu</i> - <i>Leporinus macrocephalus</i> Garavello & Britski, 1987 <i>Piava quatro-pintas</i> - <i>Schizodon intermedius</i> Garavello and Britski, 1990 <i>Amboré</i> - <i>Schizodon nasutus</i> Kner, 1858 <i>Piranha amarela</i> - <i>Serrasalmus spilopleura</i> Kner, 1858 <i>Piau verdadeiro</i> - <i>Leporinus elongatus</i> Valenciennes, 1850 <i>Piau três-pintas</i> - <i>Leporinus friderici</i> Bloch, 1794 <i>Mandi bicudo</i> - <i>Iheringichthys labrosus</i> Lütken, 1874 <i>Curimbatá</i> - <i>Prochitodus lineatus</i> Valenciennes, 1847 <i>Tambacu</i> - <i>C. macropomum</i> (fêmea) x <i>P. mesopotamicus</i> <i>Bagre africano</i> - <i>Clarias gariepinus</i> Burchell, 1822 <i>Tilápia do Nilo</i> - <i>Oreochromis niloticus</i> Linnaeus, 1758 <i>Matrinxã</i> - <i>Brycon cephalus</i> Gunther, 1869 <i>Pintado</i> - <i>Pseudoplatystoma corruscans</i> Spix and Agassiz, 1829	GABRIELLI; ORSI (2000)
Paraná	municipality of Londrina	<i>Piava quatro-pintas</i> - <i>Schizodon intermedius</i> Garavello and Britski, 1990	SILVA-SOUZA; ALMEIDA; MACHADO (2000)
Paraná	municipality of Laranjeiras do Sul	<i>Jundiá</i> - <i>Rhamdia quelen</i> Quoy & Gaimardm, 1824 <i>Carpa comum</i> - <i>Cyprinus carpio</i> Linnaeus, 1758 <i>Piau</i> - <i>Leporinus</i> sp. <i>Tambacu</i> - (<i>C. macropomum</i> (fêmea) x <i>P. mesopotamicus</i>)	CARRIEL (2014)
Paraná	municipality of Cornélio Procópio	<i>Piraputangas</i> - <i>Brycon</i> sp.	VALENTIM et al. (2003)
Rio de Janeiro	municipality of Itaocara	<i>Piabinha</i> - <i>Brycon insignis</i> Steindachner, 1876	FERNANDES et al. (2006)
São Paulo	municipality of Franca	<i>Pacu</i> - <i>Piaractus mesopotamicus</i> Holmberg, 1886 <i>Tambacu</i> (<i>C. macropomum</i> (fêmea) x <i>P. mesopotamicus</i>)	TAVARES-DIAS et al. (2001)
São Paulo	municipality of Franca	<i>Piauçu</i> - <i>Leporinus macrocephalus</i> Garavello & Britski, 1988 <i>Pacu</i> - <i>Piaractus mesopotamicus</i> Holmberg, 1887	TAVARES-DIAS et al. (1999)

(Continue)

(Continuation)

Province/ State	Location (Original name in portuguese)	Popular and scientific name Hosts	References
São Paulo	municipality of Guariba	<i>Piauçu - Leporinus macrocephalus</i> Garavello & Britski, 1987 <i>Pacu – Piaractus mesopotamicus</i> Holmberg, 1887 <i>Carpa comum - Cyprinus</i> <i>carpio</i> Linnaeus, 1758 <i>Tambacu - (C. macropomum</i> <i>(fêmea) x P. mesopotamicus)</i>	SCHALCH; MORAES (2005)
São Paulo	municipality of Jaboticabal	<i>Pacu - Piaractus mesopotamicus</i> Holmberg, 1887	MARTINS et al. (2002)
São Paulo	Taguari river, municipality of Taquarituba	<i>Lambari - Piabarchus stramineus</i> Eigenmann, 1908	NARCISO et al. (2019)
São Paulo	Taguari river, municipality of Taquarituba	<i>Saguiuri curto - Steindachnerina</i> <i>insculpta</i> Fernandez– Yopez, 1948	ACOSTA et al. (2013)
São Paulo	municipality of Pirassununga	<i>Tambaqui - Colossoma</i> <i>macropomum</i> Cuvier, 1818	BASTOS; SÃO CLEMENTE; LIMA (1996)
Santa Catarina	municipality of Florianópolis	<i>Plati - Xiphophorus maculatus</i> Günther, 1866 <i>Molinésia negra - Poecilia</i> <i>sphenops</i> Valenciennes, 1846	PIAZZA et al. (2006)
Santa Catarina	municipality of Joinville, Blumenau e Ituporanga	<i>Carpa comum - Cyprinus</i> <i>carpio</i> Linnaeus, 1758	GHIRALDELLI et al. (2006)
Santa Catarina	municipality of Itajaí	<i>Jundiá - Rhamdia quelen</i> Quoy & Gaimardm, 1824	FURTADO et al. (2019)
Sergipe	municipality of Neopólis and Propriá	<i>Lambari - Astyanax bimaculatus</i> Linnaeus, 1758	VASCONCELOS et al. (2013)
Sergipe	municipality of Aracaju	<i>Peixe japonês - Carassius</i> <i>auratus</i> Linnaeus, 1758 <i>Cascudo- hypostomus sp.</i> <i>Molinesia negra - Poecilia</i> <i>sphenops</i> Valenciennes, 1846 <i>Guppy - Poecilia reticulata</i> Peters 1859 <i>Cauda de espada - Xiphophorus</i> <i>hellerii</i> Haekel 1848	ASSIS; CAVALCANTE; BRITO (2014)
Sergipe	municipality of Propriá (San Francisco river)	<i>Tambaqui – Colossoma macropomum</i>	FUJIMOTO et al. (2019)
Mato Grosso do Sul	Great Dourados region	<i>Patinga – (Piaractus mesopotamicus x</i> <i>Piaractus brachypomus</i> Cuvier 1817) <i>Tambacu - (C. macropomum</i> <i>(fêmea) x P. mesopotamicus)</i>	ISHIKAWA et al. (2016)
Mato Grosso do Sul	Great Dourados region	<i>Tambacu - (C. macropomum</i> <i>(fêmea) x P. mesopotamicus)</i>	PEREIRA (2015)

4 Conclusion

In conclusion, this research on the parasite fauna of *Betta splendens* is the first report conducted on an *Lernaea cyprinacea* was detected on the narines, soft parts the dorsal and anal fin of this host. This new occurrence and the review presented on the previous records of the parasite in Brazil will provide the basis for further scientific studies on parasites fish tropical.

ACKNOWLEDGMENT

This study was supported by FAPEAL (Fundation Research of Alagoas State, Brazil).

REFERENCES

ACOSTA, A.A.; QUEIROZ, J.; BRANDÃO, H.; CARVALHO, E. D.; DA SILVA, R. J. Helminths of *Steindachnerina insculpta* in two distinct stretches of

the Taquari River, state of São Paulo, Brazil. **Revista Brasileira de Parasitologia Veterinária**, v. 22, n. 4, p. 539–547. 2013. <http://dx.doi.org/10.1590/S1984-29612013000400014>. Disponível em: http://www.scielo.br/scielo.php?script=sci_abstract&pid=S1984-29612013000400539&lng=en&nrm=iso. Acesso em: 21 dez. 2019.

AGOSTINHO, A. A.; GOMES, L. C.; PELICICE F. M. **Ecologia e manejo de recursos pesqueiros em reservatórios do Brasil**. Maringá, Brasil.: Eduem, 2007. 501p.

ASSIS, D. A. S.; CAVALCANTE, S. S.; BRITO, M. F. G. Aquarium trade as a potential disseminator of non-native invertebrates in Northeastern Brazil. **Neotropical Biology and Conservation**, v. 9, n. 2, p. 115–119. 2014. doi: 10.4013/nbc.2014.92.06. Disponível em: <http://revistas.unisinos.br/index.php/neotropical/article/view/nbc.2014.92.06/4223>. Acesso em 30 dez. 2019.

BARSON, M.; MULONGA, A.; NHIWATIWA, T. Investigation of a parasitic outbreak of *Lernaea cyprinacea* Linnaeus, 1758 (Crustacea: Copepoda) in fish from Zimbabwe. **African Zoology**, v. 43, n. 2, p. 175–183, 2008. <https://doi.org/10.1080/15627020.2008.11657234>. Disponível em: https://www.tandfonline.com/doi/abs/10.1080/15627020.2008.11657234?casa_token=i9i9PclG1jkAAAAA:ibSt_ZzG6WeVVGZJZopWyqdsLhd90O0gXTZFFbW3A_S680sEN4BrXrrp0Hje6Hw0n3aTEwo0-3isFr0. Acesso em 31 dez. 2019.

BASTOS, P. A. M. B.; SÃO CLEMENTE, S. C.; LIMA, F. C. Aspectos anatomo-patológicos da parasitose por *Lernaea cyprinacea* (L.). (Crustacea: Copepoda) em tambaqui (*Colossoma macropomum* Cuvier, 1818). **Revista Brasileira de Ciências Veterinárias**, v. 3, n. 1, p. 15-21, 1996. <https://doi.org/10.4322/rbcv.2015.037>. Disponível em: https://www.researchgate.net/profile/Sergio_Clemente/publication/276300549_Aspectos_anatomo-patologicos_da_parasitose_por_Lernaea_cyprinacea_L_Crustacea_Copepoda_em_tambaqui_Colossoma_macropomum_Cuvier_1818/links/568fbef908aef987e56b6373/Aspectos-anatomo-patologicos-da-parasitose-por-Lernaea-cyprinacea-L-Crustacea-Copepoda-em-tambaqui-Colossoma-macropomum-Cuvier-1818.pdf. Acesso em: 02 jan. 2020.

BERNARDINO, M. G. S.; SILVA, E. G.; BEZERRA, T. I. C.; LUCENA, R. B.; SATAKE, S. Ectoparasitologic, hematologic and histopathologic assessment of *Hoplias malabaricus* Bloch, 1794 from ponds located in Sumé municipality, state of Paraíba, Brazil. **Pesquisa Veterinária Brasileira**. v. 36, n. 7, p. 581-586. 2016.

DOI: 10.1590/S0100-736X2016000700003. Disponível em: <http://www.scielo.br/pdf/pvb/v36n7/1678-5150-pvb-36-07-00581.pdf>. Acesso em 20 set. 2019.

BERTUCCI, F.; MATOS, R. J.; DABELSTEEN, T. Knowing your audience affects male-male interactions in Siamese fighting fish (*Betta splendens*). **Animal Cognition** v. 17, n. 2, p. 229–236. 2013. DOI 10.1007/s10071-013-0655-6. Disponível em: <https://link.springer.com/content/pdf/10.1007%2Fs10071-013-0655-6.pdf>. Acesso em: 01 jan. 2020.

CARRIEL, K. C. **Levantamento da presença de Lernaea spp. (Lernaeoidea: Lernaeidae) em pisciculturas da associação de produtores de peixes de Laranjeiras do Sul-PR (Peixelar)**. 2014. 57f. Trabalho de conclusão de curso (Eng. de aquicultura) – Universidade Federal da Fronteira Sul, Laranjeiras do Sul, 2018. Disponível em: <https://rd.uffs.edu.br/bitstream/prefix/181/1/CARRIEL.pdf>. Acesso em: 20 dez. 2019.

DEMAREE JR, R. S. Ecology and external morphology of *Lernaea cyprinacea*. **American Midland Naturalist**, v. 78, n. 2, p. 416-427. 1967. Disponível em: <https://www.jstor.org/stable/pdf/2485239.pdf?refreqid=excelsior%3A4a3ff122f8631bbe4f555cfd041b2a70>. Acesso em 20 dez. 2019.

DÜPONT, A.; LOBO, E. A. First record of the occurrence of *Lernaea cyprinacea* (Linnaeus, 1758) on *Astyanax fasciatus* (Cuvier, 1819), Pardinho River, RS, Brazil. **Caderno de Pesquisa Série Biológica**. V. 23, n. 3, p. 20-25. 2011. ISSN: 1677-5600. Disponível em: <https://www.cabdirect.org/cabdirect/abstract/20123148330>. Acesso em 02 jan. 2020.

FERNANDES, G. Q.; PEREIRA, M. A. V. D. C.; MATTOS, D. G. D.; SOUZA, G. Survey of parasites in intrapopulation of cultured *Brycon insignis* Steindachner, 1876 (Pisces Characidae) from the north fluminense region, Rio de Janeiro, Brazil. **Ciência Animal Brasileira**. v. 7, n. 3, p. 309-313. 2006. ISSN 1518-2797. Disponível em: <https://www.revistas.ufg.br/vet/article/view/407/382>. Acesso em 08 dez. 2019.

FORTES, E.; HOFFMANN, R.P.; SCARIOT, J. *Lernaea cyprinacea* (Linnaeus, 1758) (Crustacea, Copepoda) parasitando peixes de água doce da Grande Porto Alegre, RS, Brasil. **Revista Brasileira de Medicina Veterinária**, v. 20, n. 2, p. 64-65. 1998. ISSN: 0100-2430. Disponível em: <https://lume.ufrgs.br/handle/10183/105389>. Acesso em 01 jan. 2020.

FURTADO, W. E.; CARDOSO, L.; FIGUEREDO, A. B.; MARCHIORI, N. C.; MARTINS, M. L. Histological and hematological alterations of silver catfish *Rhamdia quelen* highly parasitized by *Lernaea cyprinacea*. **Diseases of Aquatic Organisms**, v. 135, p. 157–168. DOI: <https://doi.org/10.3354/dao03386>. Disponível em: <https://www.int-res.com/articles/dao2019/135/d135p157.pdf>. Acesso em 02 jan. 2020.

GALLIO, M.; SILVA, A. S.; MONTEIRO, S. G. Parasitism by *Lernaea cyprinacea* in *Astyanax bimaculatus* from a dam in the municipality of Antonio Prado, Rio Grande do Sul. **Acta Scientiae Veterinariae**, 35, n. 2, p. 209–212. 2007. ISSN: 1678-0345. Disponível em: <https://www.redalyc.org/pdf/2890/289021845011.pdf>. Acesso em 21 dez. 2019.

GABRIELLI, M. A.; ORSI, M. L. Dispersão de *Lernaea cyprinacea* (Linnaeus) (Crustacea, Copepoda) na região norte do estado do Paraná, Brasil. **Revista Brasileira de Zoologia**, v. 17, n. 2, p. 395–399. 2000. <http://dx.doi.org/10.1590/S0101-81752000000200010>. Disponível em: <http://www.scielo.br/pdf/rbzool/v17n2/v17n2a10.pdf>. Acesso em: 30 dez. 2019.

GERVASONI, S. H.; CHEMES, S. B.; SCAGLIONE, M. C.; CERUTTI, R.D. First report of *Lernaea cyprinacea* (Crustacea: Lernaeidae) parasiting *Rhamdia quelen* (Pisces: Heptapteridae) in Santa Fe (Argentina) under hatchery conditions. **Revista Colombiana de Ciencias Pecuarias**, v. 31, n. 3, p. 229–234. 2018. <http://dx.doi.org/10.17533/udea.rccp.v31n3a08>. disponível em: <http://www.scielo.org.co/pdf/rccp/v31n3/0120-0690-rccp-31-03-00229.pdf>. Acesso em 31 dez. 2019.

GHIRALDELLI, L.; MARTINS, L. M.; YAMASHITA, M. M.; JERONIMO, G. T. Ectoparasites on the haematological parameters of Nile tilapia and carp culture in the State of Santa Catarina South Brazil. **Journal of Fisheries and**

Aquatic Science, v. 1, n. 3, p. 270–276. 2006. DOI: 10.3923/jfas.2006.270.276. Disponível em: <https://scialert.net/fulltextmobile/?doi=jfas.2006.270.276>. Acesso 09 jan. 2020.

GIANNECCHINI, L.G.; MASSAGO, H.; FERNANDES, J. B. K. Effects of photoperiod on reproduction of Siamese fighting fish *Betta splendens*. **Revista Brasileira de Zootecnia**, v. 41, n. 4, p. 821–826. 2012. <http://dx.doi.org/10.1590/S1516-35982012000400001>. Disponível em: <http://www.scielo.br/pdf/rbz/v41n4/01.pdf>. Acesso em 31 dez. 2019.

FABREGAT, T. E.; WOSNIAK, B.; TAKATA, R.; MIRANDA-FILHO, K. C.; FERNANDES, J. B.; PORTELLA, M. C. Larviculture of siamese fighting fish *Betta splendens* in low-salinity water. **Boletim do Instituto de Pesca**, v. 43, n. 2, p. 164–171, 2017. Doi: 10.20950/1678-2305.2017v43n2p164. Disponível em: <https://www.pesca.sp.gov.br/Artigo-2-BIP-086.pdf>. Acesso em 31 dez. 2019.

FUJIMOTO, R. Y.; HIDE, D. M. V.; PAIXÃO, P. E. G.; ABE, H. A.; DIAS, J. A. R.; SOUSA, N. C.; COUTO, M. V. S.; SILVA, R. V. B.; MADI, R. R.; BENAVIDES, M. V.; ISHIKAWA, M. M.; CHAGAS, E. C.; BOIJINK, C. L.; DOMPIERI, M. H. G.; PEREIRA, A. M. L.; MACIEL, P. O. Fauna parasitária e relação parasito-hospedeiro de tambaquis criados na região do Baixo São Francisco, nordeste do Brasil. **Arquivo Brasileiro de Medicina Veterinária e Zootecnia**, v. 71, n. 2, 2019. <http://dx.doi.org/10.1590/1678-4162-10306>. Disponível em: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0102-09352019000200563. Acesso em 02 jan. 2020.

HOSSAIN, M. M. M.; FERDOUSHI, J.; RUPON, A. H. Biology of anchor worms (*Lernaea cyprinacea*). **Journal of Entomology and Zoology Studies**, v. 6, n. 1, p. 910–917, 2018. ISSN: 2349-6800. Disponível em: https://www.researchgate.net/profile/Mmm_Hossain/publication/327652082_Biology_of_anchor_worms_Lernaea_cyprinacea/links/5b9bb5fe458515747c92fba/Biology-of-anchor-worms-Lernaea-cyprinacea.pdf. Acesso em: 06 nov. 2019.

INNAL, D.; AVENANT-OLDEWAGE, A.; DOGANGIL, B.; STAVRESCU-BEDIVAN, M. M.; OZMEN, O.; MAVRUK, S. Susceptibility of endemic and non-indigenous fish to *Lernaea cyprinacea* (Copepoda: Lernaeidae): a case study from Düger Spring Creek (Burdur-Turkey). **Bulletin of the European Association of Fish Pathologists**, v. 37, n. 3, p. 100–109, 2017. ISSN: 0108-0288. Disponível em: https://www.researchgate.net/profile/MALA_STAVRESCU_BEDIVAN/publication/317592477_Susceptibility_of_endemic_and_non-indigenous_fish_to_Lernaea_cyprinacea_Copepoda_Lernaeidae_a_case_study_from_Duger_Spring_Turkey/links/594bdbcf458515e70348936c/Susceptibility-of-endemic-and-non-indigenous-fish-to-Lernaea-cyprinacea-Copepoda-Lernaeidae-a-case-study-from-Dueger-Spring-Turkey.pdf. Acesso em: 02 jan. 2020.

ISHIKAWA, M. M.; JERÔNIMO, G. T.; VENTURA, A. S.; PEREIRA, N. L.; SILVA, T. S. C.; ZANON, R. B.; FUJIMOTO, R. Y.; CHAGAS, E. C.; MACIEL, P. O.; BENAVIDES, M. V.; MARTINS, M. L. **Parasitos de peixes redondos produzidos na região**

da Grande Dourados - MS: características e possíveis soluções.

Jaguariúna: Embrapa Meio Ambiente, 2016. 40 p. Disponível em: <https://ainfo.cnptia.embrapa.br/digital/bitstream/item/158116/1/2016DC08.pdf>. Acesso em 30 out. 2019.

KOYUN, M.; ULUPINAR, M.; MART, A.

First Record of *Lernaea cyprinacea* L. 1758 (Copepoda: Cyclopoida) on *Cyprinion macrostomus* Heckel, 1843 from Eastern Anatolia, Turkey. **Biharean Biologist**, v. 9, n. 1, p. 44-46. 2015. ISSN: 1843-5637. Disponível em: http://biozoojournals.ro/bihbiol/cont/v9n1/bb_141130_Koyun.pdf. Acesso em: 11 dez. 2019.

KUMAR, S.; BEHERA, S.; NAGESH, T. S.; RINKU, G.; CH, B. P. Effect of 17 α -methyltestosterone on the growth performance of a fighting fish (*Betta splendens*). **Journal of Experimental Zoology, India**, v. 19, n. 1, p. 89-93, 2016. ISSN 0972-0030. Disponível em: <https://www.cabdirect.org/cabdirect/abstract/20163131593>. Acesso em: 02 jan. 2020.

LUQUE, J. L. Biologia, epidemiologia e controle de parasitos de peixes. **Revista Brasileira de Parasitologia Veterinária**, v. 13, Supl. 1, p. 161-165, 2004. ISSN 1984-2961. Disponível em: http://www.ufrjr.br/laboratorio/parasitologia/arquivos/publicacao/48_LIVRO.pdf. Acesso em: 02 jan. 2020.

MACEDA VEIGA, A.; DOMÍNGUEZ DOMÍNGUEZ, O.; ESCRIBANO ALACID, J.; LYONS, J. The aquarium hobby: Can sinners become saints in freshwater fish conservation? **Fish and Fisheries**, v. 17, n. 3, p. 860– 874, 2016. <https://doi.org/10.1111/faf.12097>. Disponível em: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/faf.12097>. Acesso em 02 jan. 2020.

MACLAREN, R. D.; WISNIEWSKI, K.; MACLAREN, C. Environmental concentrations of metformin exposure affect aggressive behavior in the Siamese fighting fish, *Betta splendens*. **Plos One**, v.13, n.5, 2018. <https://doi.org/10.1371/journal.pone.0197259>. Disponível em: <https://journals.plos.org/plosone/article/file?id=10.1371/journal.pone.0197259&type=printable>. Acesso em 02 jan. 2020.

MAGALHÃES, A. L. B. First record of lernaecosis in a native fish species from a natural environment in Minas Gerais state, Brazil. **Pan-American Journal of Aquatic Sciences**. v. 1, n. 1. p. 8-10, 2006. ISSN: 1809-9009. Disponível em: https://www.researchgate.net/publication/259738438_First_record_of_lernaecosis_in_a_

[native_fish_species_from_a_natural_environment_in_Minas_Gerais_state_Brazil](https://www.researchgate.net/publication/259738438_First_record_of_lernaecosis_in_a_native_fish_species_from_a_natural_environment_in_Minas_Gerais_state_Brazil). Acesso em 02 dez. 2019.

MANCINI, M.; RODRIGUEZ, C; ORTIZ, M.; SALINAS, V.; TANZOLA, R. Lernaecosis en peces silvestres y cultivados del centro de Argentina. **Biología Acuática**, v. 24, p. 33-41, 2008. ISSN: 0326-1638. Disponível em: http://sedici.unlp.edu.ar/bitstream/handle/10915/67980/Documento_completo.pdf-PDFA.pdf?sequence=1&isAllowed=y. Acesso em 02 jan. 2020.

MARTINS, M. L.; ONAKA, E. M.; MORAES, F. R.; BOZZO, F. R.; MELLO, A.; PAIVA, F. C.; GONÇALVES, A. Recent studies on parasitic infections of freshwater cultivated fish in the State of São Paulo, Brazil. **Acta Scientiarum Agronomy**, V. 24, n. 4, p. 981-985, 2002. ISSN 1679-9275. Disponível em: <https://pdfs.semanticscholar.org/b80a/2e0c2af72c234a10b3904d2744cdf81a0f43.pdf>. Acesso em 27 nov. 2019.

MIRZAEI, M. Prevalence and histopathologic study of *Lernaea cyprinacea* in two species of ornamental fish (*Poecilia latipinna* and *Xiphophorus helleri*) in Kerman, South-East Iran. **Turkiye parazitolojii dergisi**. v. 39, n. 3, p. 222-226. 2015. DOI: 10.5152/tpd.2015.3960. Disponível em: <https://www.ncbi.nlm.nih.gov/pubmed/26470930> Acesso em 02 jan. 2019.

MOTLAGH, S. P.; ZAREJABAD, A. M.; NASRABADI, R. G.; AHMADIFAR, E.; MOLAEI, M. Haematology, morphology and blood cells characteristics of male and female Siamese fighting fish (*Betta splendens*). **Comparative Clinical Pathology**, v. 21, n. 1, p. 15–21, 2012. doi:10.1007/s00580-010-1058-6. Disponível em: <https://link.springer.com/article/10.1007/s00580-010-1058-6>. Acesso em 02 dez. 2019.

NARCISO, R. B.; ACOSTA, A. A.; NOBILE, A. B.; LIMA, N. P.; FREITAS-SOUZA, D.; SILVA, R. J. *Lernaea cyprinacea* (Copepoda: Lernaecidae) in *Piabarchus stramineus* (Characiformes: Characidae) from the Taquari River, São Paulo State, Brazil. **Biologia**, v.74, n. 9, p. 1171–1179, 2019. <http://dx.doi.org/10.2478/s11756-019-00240-z>. Disponível em: <https://link.springer.com/article/10.2478/s11756-019-00240-z>. Acesso em 02 jan. 2020.

PENNING, M.; REID, G. M. G.; KOLDEWEY, H.; DICK, G.; ANDREWS, B.; ARAI, K. In P. GARRATT, S.; GENDRON, J.; LANGE, K.; TANNER, S.; TONGE, P.; SANDE, P.; WARMOLTS, D.; GIBSON, C. (Eds.), **Turning the Tide: A Global Aquarium Strategy for Conservation and Sustainability**. Bern: World Association of Zoos and Aquariums. 2009. Disponível

em: <https://www.waza.org/wp-content/uploads/2019/03/Aquarium-strategy-EN.pdf>. Acesso em: 02 jan. 2020.

PEREIRA, N. L. **Parasitas metazoários de pacu (*Piaractus mesopotamicus*) e dos híbridos patinga e tambacu cultivados na região da grande Dourados-MS, Brasil**. 2015. 38f. Dissertação (Mestrado em desenvolvimento sustentável) - Universidade Federal do Paraná, Palotina, 2015. Disponível em: <https://acervodigital.ufpr.br/bitstream/handle/1884/40969/R%20-%20D%20-%20NATHALIA%20LOPEZ%20PEREIRA.pdf?sequence=2&isAllowed=y>. Acesso em 02 jan. 2020.

PIAZZA, R. S.; MARTINS, M. L.; GUIRALDELLI, L.; YAMASHITA, M. M. Parasitic diseases of freshwater ornamental fishes commercialized in Florianópolis, Santa Catarina, Brazil. **Boletim do Instituto Pesca**. v. 32, p. 51-57. 2006. ISSN 0046-9939. Disponível em: https://www.pesca.sp.gov.br/32_1_51-57.pdf. Acesso em: 29 dez. 2019.

PLEEGING, C.C.F.; MOONS, C.P.H. Potential welfare issues of the Siamese fighting fish (*Betta splendens*) at the retailer and in the hobbyist Aquarium. **Vlaams Diergeneeskundig Tijdschrift**, v. 86, p. 213-223, 2017. ISSN: 0303 9021. Disponível em: <https://biblio.ugent.be/publication/8533462/file/8605425.pdf>. Acesso em: 29 dez. 2019.

QUEROL, M. V. M.; QUEROL, E.; PESSANO, E. F. C.; AZEVEDO, C. L. O. Ocorrência da carpa húngara, *Cyprinus carpio* (Linnaeus, 1758) e disseminação parasitária, no Arroio Felizardo, Bacia do médio Uruguai, Uruguaiana, RS, Brasil. **Biodiversidade Pampeana**. v. 3, n. 1, p. 21-23. 2005. ISSN 1679-6179. Disponível em: <http://revistaseletronicas.pucrs.br/ojs/index.php/biodiversidadepampeana/article/view/2535/1982>. Acesso em 20 dez. 2019.

SALINAS, Z. A.; BABINI, M. S.; GRENAT, P. R.; BIOLÉ, F. G.; MARTINO, A. L.; SALAS, N. E. Effect of parasitism of *Lernaea cyprinacea* on tadpoles of the invasive species *Lithobates catesbeianus*. **Heliyon**, v. 5, n. 6, e01834, 2019. <https://doi.org/10.1016/j.heliyon.2019.e01834>. Disponível em: <https://reader.elsevier.com/reader/sd/pii/S2405844018355804?token=0CFC0DA6B467DE8A1332CBA143FDC999D0806C37878AF8D204FC405E986B03F9082AB8C850E08A58915413656AB581C8>. Acesso em 02 jan. 2020.

SANTOS, M.; BRASIL-SATO, M. C. Parasitic community of *Franciscodoras marmoratus* (Reinhardt, 1874) (Pisces: Siluriformes: Doradidae) from the Upper São Francisco River, Brazil. **Brazilian Journal of Biology**. V. 66, n. 3, p. 931-938. 2006. <http://dx.doi.org/10.1590/S1519-69842006000500019>. Disponível em: <http://www.scielo.br/pdf/bjb/v66n3/18.pdf>. Acesso em 02 dez. 2019.

SANTOS, E. L.; SILVA, F. C. B.; PONTES, E. C., LIRA, R. C.; CAVALCANTI, M. C. A. Residue of the process red propolis extract in commercial ration on performance of Nile tilapia fingerlings (*Oreochromis niloticus*). **Comunicata Scientiae**. V. 4, n. 2, p. 179-185. 2013. DOI: <https://doi.org/10.14295/cs.v4i2.261>. Disponível em: <https://www.comunicatascientiae.com.br/comunicata/article/view/261/178>. Acesso em: 18 dez. 2019.

SANTOS, M.A.; JERÔNIMO, G.T.; CARDOSO, L.; TANCREDO, K.R.; MEDEIROS, P.B.; FERRAREZI J.V.; GONÇALVES, E.L.T.; ASSIS, G.C.; MARTINS, M.L. Parasitic fauna and histopathology of farmed freshwater ornamental fish in Brazil. **Aquaculture**. v. 470, p.103-109. 2017. <https://doi.org/10.1016/j.aquaculture.2016.12.032>. Disponível em: <https://www.sciencedirect.com/science/article/abs/pii/S0044848616309048>. Acesso em 27 abr. 2020.

SANTOS, E. L.; LIRA, R. C.; SOUZA, C. A.; SANTOS, I. V. V. S.; SILVA, F. C. B.; ALBUQUERQUE, I. C. M. Desempenho de *Betta splendens* Associados a Diferentes Frequências Alimentares. **Revista Científica de produção Animal**. V. 2014. DOI: <http://dx.doi.org/10.15528/2176-4158/rcpa.v16n1p10-16>. Disponível em: <https://pdfs.semanticscholar.org/1417/12806683e69c5b3cd3f996d824e7b534ed9c.pdf>. Acesso em 12 dez. 2019.

SANTOS, E. L.; SILVA, L. L. A.; SOARES, E. C.; OLIVEIRA, W. D. S.; LIMA, M. R.; OLIVEIRA, J. M.; SANTOS, E. L. Frequência e horário de arraçoamento sobre o desempenho de machos *Betta splendens*. **Biota Amazônia**, v. 9, n. 3, p. 30-34, 2019. DOI:<http://dx.doi.org/10.18561/21795746/biotaamazonia.v9n3p30-34>. Disponível em: <https://periodicos.unifap.br/index.php/biota/article/view/4027/v9n3p30-34.pdf>. Acesso em: 01 jan. 2020.

SALES, A. D.; VERAS, G. C.; BRABO, M. F.; ABRUNHOSA, F. A.; ALVES, A. X.; DIAS, B. C. B.; CAMPELO, D. A. V.; FERREIRA, M. S. S. Fotoperíodo e frequência alimentar na larvicultura do peixe beta. **Arquivo Brasileiro de Medicina Veterinária e Zootecnia**. v. 68, n. 4, p. 1062-1068. 2016. <http://dx.doi.org/10.1590/1678-4162-8746>. Disponível em:

<http://www.scielo.br/pdf/abmvz/v68n4/0102-0935-abmvz-68-04-01062.pdf>. Acesso em: 30 out. 2019.

SCHALCH, S. H. C.; MORAES, F. R. Seasonal distribution of gill parasites in fishes from feefishing at Guariba, São Paulo State, Brazil. **Revista Brasileira de Parasitologia Veterinária**, v. 14, p. 141–146, 2005. ISSN: 0103-846X. Disponível em: https://www.researchgate.net/publication/7327793_Seasonal_distribution_of_gill_parasites_in_fishes_from_feefishing_at_Guariba_Sao_Paulo_State_Brazil. Acesso em: 02 jan. 2020.

SILVA-SOUZA, A. T.; ALMEIDA, S. C.; MACHADO, P. M. Effect of the infestation by *Lernaea cyprinacea* Linnaeus, 1758 (Copepoda, Lernaeidae) on the leucocytes of *Schizodon intermedius* Garavello and Britski, 1990 (Osteichthyes, Anostomidae). **Revista Brasileira de Biologia**, v. 60, n. 2, p. 217–220. 2000. <http://dx.doi.org/10.1590/S0034-71082000000200004>. Disponível em: <http://www.scielo.br/pdf/rbbio/v60n2/a04v60n2.pdf>. Acesso em: 09 dez. 2019.

SOARES, I. A.; SALINAS, V.; PONTI, O. D.; MANCINI, N. A.; LUQUE, J. L. First molecular data for *Lernaea cyprinacea* (Copepoda: Cyclopoida) infesting *Odontesthes bonariensis*, a commercially important freshwater fish in Argentina. **Brazilian Journal Veterinary Parasitology**, v. 27, n. 1, p. 106–109, 2018. <http://dx.doi.org/10.1590/s1984-29612018005>. Disponível em: <http://www.scielo.br/pdf/rbvp/2018nahead/1984-2961-rbvp-S1984-29612018005.pdf>. Acesos em 30 set. 2019.

TAVARES-DIAS, M.; SCHALCH, S. H. C.; MARTINS, M. L.; SILVA, E. D.; MORAES, F. R.; PERECIN, D. Haematology of parasitized Brazilian teleost fishes. I. Parameters of *Leporinus macrocephalus* Garavelo and Britski, 1988 (Anostomidae) and *Piaractus mesopotamicus* Holmberg, 1887 (Characidae). **Acta Scientiarum Biological Science**, v. 21, n. 3, p. 337–342. 1999. ISSN: 1415-6814. Disponível em: <http://periodicos.uem.br/ojs/index.php/ActaSciBiolSci/article/view/4440/3026>. Acesso em: 23 dez. 2019.

TAVARES-DIAS, M.; MARTINS, M. L.; MORAES, F. R.; KRONKA, S. Fauna parasitária de peixes oriundos de “pesque-pague” do município de Franca, São Paulo, Brasil I Protozoários. **Revista Brasileira de Zoologia**, v. 18, p.67–79, 2001. <http://dx.doi.org/10.1590/S0101-81752001000500006>. Disponível em: <https://repositorio.unesp.br/bitstream/handle/11449/3147/S0101-81752001000500006.pdf?sequence=1&isAllowed=y>. acesso em: 02 jan. 2020.

TAVARES-DIAS, M.; DIAS-JÚNIOR, M. B. F.; FLORENTINO, A. C.; SILVA, L. M. A.; CUNHA, A. C.

Distribution pattern of crustacean ectoparasites of freshwater fish from Brazil. **Revista Brasileira de Parasitologia Veterinária**, v. 24, n. 2, p. 136–147, 2015; <http://dx.doi.org/10.1590/S1984-29612015036>. Disponível em: <http://www.scielo.br/pdf/rbvp/v24n2/1984-2961-rbvp-24-2-136.pdf>. Acesso em: 09 fev. 2019.

THATCHER, V. E. **Amazon Fish Parasites**. In: ADIS J, ARIAS JR, RUEDA-DELGADO G, WANTZEN KM, editors. Aquatic Biodiversity in Latin America. 2nd ed. Sofia-Moscow: Pensoft Publishers; 2006. 508p.

TORRES-VALENCIA, G. A.; IMUES-FIGUEROA, M. A.; BUCHELI-FUELANTALA, J. F. Larvicultura del *Betta splendens* utilizando rotíferos *Brachionus calyciflorus* e infusorios como fuente de alimento. **Orinoquia**, v. 20, n. 2, p. 71–77, 2016. ISSN: 0121-3709. Disponível em: <https://www.redalyc.org/pdf/896/89659214008.pdf>. Acesso em: 21 set. 2019.

THONGPRAJUKAEW, K.; KOVITVADHI, U.; KOVITVADHI, S.; SOMSUEB, P.; RUNGRUANGSAK-TORRISEN, K. Effects of different modified diets on growth, digestive enzyme activities and muscle compositions in juvenile Siamese fighting fish (*Betta splendens* Regan, 1910). **Aquaculture**, v. 322–323, n. 21, p. 1–9. 2011. <https://doi.org/10.1016/j.aquaculture.2011.10.006>. Disponível em: <https://www.sciencedirect.com/science/article/abs/pii/S0044848611007964>. Acesso em: 02 jan. 2020.

VASCONCELOS, A. C. P.; LOPES, A. C. M.; SANTOS, J. M. S.; JERALDO, V. L. S.; MELO, C.M.; MADI, R. R. Molecular analysis and biodiversity of metazoan parasites of the yellow tail lambari, *Astyanax aff. bimaculatus* (Teleostei, Characidae), in lower San Francisco, northeastern Brazil. **Neotropical Helminthology**, v. 7, n. 1, p. 41–49, 2013. ISSN-e: 1995-1043. Disponível em: <https://dialnet.unirioja.es/servlet/articulo?codigo=4435932>. Acesso em 02 jan. 2020.

VALENTIM, M.; VARGAS, L.; MOREIRA, H.M.; RIBEIRO, R.P. Comparison of DNA extraction protocols in *Lernaea* sp. (Copepoda: Cyclopoida). **Acta Scientiarum Animal Science**, v. 25, n. 2, p. 219–222. 2003. DOI: 10.4025/actascianimsci.v25i2.1973. Disponível em: https://www.researchgate.net/publication/237979592_Comparacao_de_protocolos_para_extracao_do_DNA_de_Lernaea_sp_Copepoda_Cyclopoida_-_DOI_104025actascianimsciv25i21973. Acesso em 13 dez. 2019.

VASAGAN, K. P. K.; RAJKUMAR, M.; JEAN-PAUL, T.; BALASUBRAMANIAN, T. Note on *Lernaea Cyprinacea* (Crustacea, Copepoda, Lernaeidae)

Parasitizing the Cultured Sailfin Molly *Poecilia latipinna* and Their Control with Salinity Treatment. **Journal of Fisheries and Aquatic Science**, v.1, n. 3, p. 284-290, 2006. ISSN: 1816-4927. Disponível em: <http://docsdrive.com/pdfs/academicjournals/jfas/2006/284-290.pdf>. Acesso em: 02 jan. 2020.

WAICHEIM, M. A.; BLASETTI, G.; CORDERO, P.; RAUQUE, C. A.; VIOZZI, G. P. The invasive copepod *Lernaea cyprinacea* Linnaeus, 1758 (Copepoda, Cyclopoida, Lernaeidae): first record for Neuquén River, Patagonia, Argentina. **Check List**. v. 13, n. 6, p. 997–1001. 2017. <https://doi.org/10.15560/13.6.997>. Disponível em: <https://checklist.pensoft.net/article/20205/>. Acesso em 07 set. 2019.

ZUANON, J. A. S.; SALARO, A. L.; FURUYA, W. M. Produção e nutrição de peixes ornamentais. **Revista Brasileira de Zootecnia**, 40, n. 1, p. 165-174. 2011. ISSN 1806-9290. Disponível em: <http://www.sbz.org.br/revista/artigos/66271.pdf>. Acesso em 09 nov. 2019.